



# Pharmacodynamic evaluation of the Intracellular activity of CEM-101, a novel fluoro ketolide, towards *Staphylococcus aureus*, *Listeria monocytogenes*, and *Legionella pneumophila* in human THP-1 macrophages.

Sandrine Lemaire, Françoise Van Bambeke, Paul M. Tulkens

Unité de pharmacologie cellulaire et moléculaire & Louvain Drug Research Institute, Brussels, Belgium.

Mailing address:

P.M. Tulkens  
UCL 73.70 av. Mounier 73,  
1200 Brussels – Belgium,  
tulkens@facm.ucl.ac.be



PXXX

## Abstract

### Background:

Macrolides accumulate in eukaryotic cells and are considered advantageous for the treatment of intracellular infections. Ketolides recover activity against erythromycin-resistant organisms. CEM-101, a novel fluoro ketolide with a 11,12 carbamate-butyl-[1,2,3]-triazolyl-phenylamino sidechain, demonstrates enhanced potency compared to tetracyclines. We have assessed the cellular accumulation and intracellular activity of CEM-101 towards the intracellular forms of *Staphylococcus aureus*, *Listeria monocytogenes*, and *Legionella pneumophila* in comparison with azithromycin (AZM), clarithromycin (CLR), and telithromycin (TEL).

### Methods:

All experiments were performed with the human macrophage cell line THP-1. Drug accumulation was measured using bioassay. Intracellular activity measured over time and concentration by following the change in cell-associated CFU compared to post-phagocytosis levels (see details in JAC 2004,54:289-9 [*L. monocytogenes* – strain EGD]; AAC 2006, 50:841-51 [*S. aureus*; strain ATCC 25923]; similar protocol for *L. pneumophila* [strain ATCC 33153]).

### Results:

Uptake of CEM-101 was linear over time, reaching accumulation levels about 375-fold within 24 h (AZM, 160 x, CLR, 30 x, TEL, 21 x). Accumulation was suppressed by acidic pH or addition of the proton ionophore monensin, but not modified by verapamil or gemfibrozil (preferential inhibitors of P-gp and MRP, respectively). MIC and intracellular activities (developed in a concentration-dependent fashion [Hill equation] are shown in Table.

	CEM-101			AZM			CLR			TEL		
	MIC <sup>a</sup>	Cs <sup>b</sup>	E <sub>max</sub> <sup>c</sup>	MIC <sup>a</sup>	Cs <sup>b</sup>	E <sub>max</sub> <sup>c</sup>	MIC <sup>a</sup>	Cs <sup>b</sup>	E <sub>max</sub> <sup>c</sup>	MIC <sup>a</sup>	Cs <sup>b</sup>	E <sub>max</sub> <sup>c</sup>
S.a.	0.06	0.022	-0.86	0.5	> 10	0.04	0.5	0.84	-0.18	0.25	0.63	-0.29
L.m.	0.004	0.11	-0.66	1	11.6	-0.81						
L.p.	0.004	0.018	-1.03	0.016	2.90	-0.83	0.007	0.12	-0.71	0.007	0.06	-0.63

<sup>a</sup>mg/L = static concentration (mg/L) at 24 h; <sup>b</sup>μg/L; <sup>c</sup>CFU at 24h compared to the post-phagocytosis inoculum

CEM-101 caused slightly higher or similar maximal efficacy compared to AZM, CLR or TEL but considerably higher relative potency (lower EC<sub>50</sub> and C<sub>1</sub>) in relation to its lower MIC when expressed on a mass basis (differences in EC<sub>50</sub> and C<sub>1</sub> between drugs largely vanish if data are expressed as multiples of the MIC).

**Conclusions:** CEM-101 is a ketolide with enhanced cellular accumulation. It shows improved intracellular potency (on a weight basis) in comparison with AZM, CLR and TEL. In this *in vitro* model (mainly due to its larger intrinsic activity [lower MICs] against target organisms). This should lead to enhanced *in vivo* potency if using doses similar to those of the comparators tested here.

## References

- Schwartz J, et al. *Cell Culture*. 1991. Triazole derivatives. *Ann. J. Med.* 155:115-116.
- Zhang G, et al. *Antimicrob. Agents Chemother.* 2003. *In vitro* activity of a novel fluoro ketolide, CEM-101, against *Staphylococcus aureus* and *Legionella pneumophila* in THP-1 macrophages. *Antimicrob. Agents Chemother.* 2003; 47:1442-1448.
- Smith C, et al. *Antimicrob. Agents Chemother.* 2003. *In vitro* activity of a novel fluoro ketolide, CEM-101, against *Staphylococcus aureus* and *Legionella pneumophila* in THP-1 macrophages. *Antimicrob. Agents Chemother.* 2003; 47:1442-1448.
- Fernandez F, et al. *Antimicrob. Agents Chemother.* 1995. New macrolide active against *Streptococcus pyogenes* with indole and coumarine type of macrocyclic-etheramino-B-2 residues. *Antimicrob. Agents Chemother.* 1995; 39:1733-1738.
- Aguiar A, et al. *Antimicrob. Agents Chemother.* 2003. *In vitro* activity of a novel fluoro ketolide, CEM-101, against *Staphylococcus aureus* and *Legionella pneumophila* in THP-1 macrophages. *Antimicrob. Agents Chemother.* 2003; 47:1442-1448.
- Ueno-Sakai S, et al. *Antimicrob. Agents Chemother.* 2003. *In vitro* activity of a novel fluoro ketolide, CEM-101, against *Staphylococcus aureus* and *Legionella pneumophila* in THP-1 macrophages. *Antimicrob. Agents Chemother.* 2003; 47:1442-1448.
- Beck-Broichsavar M, et al. *Antimicrob. Agents Chemother.* 2003. *In vitro* activity of a novel fluoro ketolide, CEM-101, against *Staphylococcus aureus* and *Legionella pneumophila* in THP-1 macrophages. *Antimicrob. Agents Chemother.* 2003; 47:1442-1448.


## Introduction

Macrolides accumulate inside eukaryotic cells, which has been considered advantageous for the treatment of intracellular infections (1,2), even though we know that these antibiotics express only a minimal fraction of their antibacterial potential intracellularly (3).

11,12-carbamate analogs of clarithromycin carrying a lipophilic side chain show improved activity compared to the parent compound (4). Together with the removal of the cladinoside, this led to the discovery and development of the telithromycin (5), the first ketolide to reach clinical approval. CEM-101 is a novel fluoro ketolide containing an 11,12-carbamate-butyl-[1,2,3]-triazolyl-aminophenyl sidechain (see structure hereunder) that shows enhanced potency compared to telithromycin (6). We have assessed its cellular accumulation and intracellular activity using models that have been developed for the study of the intracellular pharmacodynamics of antibiotics (3,7).

### Structural formula of CEM-101

The molecule possesses a 11,12-carbamate-butyl-[1,2,3]-triazolyl-aminophenyl sidechain and a fluorine atom (green arrowhead) substituting position 2 of the macrocycle. The molecule is monocationic at neutral or moderately acidic pH, as the calculated pKa of the aminophenyl group is less than 4 (vs. ~8.5 for the desosamine).



## Methods

**Cells:** THP-1 myelomonocytic cells (ATCC TIB-202) with macrophage-like activity (7).

**Antibiotic assay:** microbiological method (7) and accumulation calculated as apparent intracellular/extracellular concentration ratio (7) based on assay of cell protein content (7)

**Bacterial strains:** *S. aureus* ATCC 25923 (methicillin-sensitive), *L. monocytogenes* strain EGD and *L. pneumophila* strain ATCC 33153

**Cell infection and assessment of antibiotic intracellular activities:** Time- (0-24h) and concentration-dependent (24 h for *S. aureus* and *L. monocytogenes*; 48h for *L. pneumophila*) experiments as described in ref. 3 (minor adaptations for *L. pneumophila*). Data from conc.-dependent experiments analyzed by non-linear regression using Hill's equation (7) to calculate pharmacological descriptors (E<sub>max</sub>: maximal reduction of the intracellular inoculum (in log<sub>10</sub> units) for an infinitely large antibiotic concentration; EC<sub>50</sub>: increase in intracellular inoculum (in log<sub>10</sub> units) for an infinitely low antibiotic concentration; EC<sub>10</sub>: antibiotic concentration yielding a response half-way between E<sub>max</sub> and E<sub>50%</sub>; C<sub>1</sub>: antibiotic concentration yielding a static effect.

## Results




Figure 1: Accumulation of CEM-101 vs. comparators in uninfected cells at 37°C (C=10 mg/L). A, influence of incubation time; B, influence of pH (30 min); C, influence of monensin (10<sup>-6</sup> M; 30 min), verapamil (P-gp inhibitor; 150 μM; 24 h) or gemfibrozil (MRP inhibitor; 250 μM; 24 h).




Figure 2: Comparative susceptibilities of *S. aureus* ATCC 25923 and *L. monocytogenes* EGD to CEM-101, telithromycin, azithromycin, and clarithromycin.

Figure 3: Short-term time-kill effect of CEM-101 and azithromycin towards *S. aureus* (ATCC 25923) in broth vs. in THP-1 macrophages.




Figure 4: Concentration-effect relationships for CEM-101, telithromycin, clarithromycin, and azithromycin towards *S. aureus* (ATCC 25923) in broth or in THP-1 macrophages. Ordinate: change in CFU (Δ log<sub>10</sub> CFU) per mL (broth) or per mg of cell protein (THP-1 macrophages) at 24 h compared to initial inoculum. Abscissa: weight concentrations (upper rows) or multiples of MIC (measured at pH 7.4; lower rows).




Figure 5: Concentration-effect relationships for CEM-101 and azithromycin towards intraphagocytic *L. monocytogenes* and *L. pneumophila*. Ordinate: change in CFU (Δ log<sub>10</sub> CFU) per mg of cell protein at 24 h (*L. monocytogenes*) or 48 h (*L. pneumophila*) compared to initial post-phagocytosis inoculum. Abscissa: weight concentrations (upper rows) or multiples of MIC (measured at pH 7.4; lower rows).

## Pharmacological parameters of the dose response of *S. aureus* to the antibiotics tested in this study (data from Figure 4)

antibiotic	broth			THP-1 macrophages		
	E <sub>max</sub>	EC <sub>50</sub>	C <sub>1</sub>	E <sub>max</sub>	EC <sub>50</sub>	C <sub>1</sub>
CEM-101	-1.37	mg/L 0.03 x MIC 0.48	0.06 0.88	-0.86	mg/L 0.0068 x MIC 0.11	0.022 0.35
telithromycin	-1.00	mg/L 0.12 x MIC 0.46	0.29 0.99	-0.29	mg/L 0.024 x MIC 0.097	0.63 1.04
azithromycin	-1.23	mg/L 1.76 x MIC 3.55	3.4 8.7	0.04	mg/L 0.11 x MIC 0.22	>50 >100
clarithromycin	-1.41	mg/L 0.80 x MIC 1.59	1.32 2.65	-0.18	mg/L 0.046 x MIC 0.093	0.84 1.68

**ACKNOWLEDGEMENTS:** We thank M.C. Cambier and C. Meson for technical assistance. This work was supported by the Belgian *Fonds de la Recherche Scientifique Médicale (FRSM)* and by a grant-in-aid from Cempira Pharmaceuticals.